



**CONTROL PROGRAM**  
for biological products  
**EXAMPLE**



# GREENHOUSE CUCUMBER

SEEDLING PRODUCTION		AFTER TRANSPLANTING			
DAMPING-OFF		ROOT DISEASES		GUMMY STEM BLIGHT <i>(Didymella)</i> GREY MOULD <i>(Botrytis)</i>	MANAGEMENT OF STRESS FACTORS
Peat mixtures: <b>MYCOSTOP®</b>	Rockwool: <b>LALSTOP® G46 WG</b>	Peat mixtures: <b>MYCOSTOP®</b> <i>Fusarium, Pythium</i>	Rockwool: <b>LALSTOP® G46 WG</b> <i>Pythium, Fusarium, Rhizoctonia, Phytophthora</i>	<b>LALSTOP® G46 WG</b>	<b>LALSTIM® OSMO</b>
Treatment immediately after sowing or germination.		First application at transplanting, repeat at intervals of 4-6 weeks.		First application after transplanting, repeat at intervals of 3 weeks.	First application 24 h before stress conditions. Repeat at intervals of 3 weeks during the stress period.
DRENCH OR SUBSTRATE SPRAY <sup>1)</sup>		DRIP IRRIGATION OR SUBSTRATE SPRAY/DRENCH <sup>1)</sup>		FOLIAR SPRAY <sup>2)</sup>	FOLIAR SPRAY <sup>2)</sup> OR DRIP IRRIGATION <sup>1)</sup> (only with rockwool substrate)
2–10 g/100 m <sup>2</sup>	50–100 g/100 m <sup>2</sup>	5–8 g/1000 plants	20–25 g/1000 plants	250–500 g/ha (Spraying volume 500–2000 l/ha)	2–3 kg/ha (Spraying volume 400–1000 l/ha)

1) The amount of water can be adjusted according to the humidity of the substrate and the need for irrigation.

2) **LALSTIM® OSMO** and **LALSTOP® G46 WG** can be sprayed on the foliage in the same mixture.

**LALSTIM® OSMO** can be applied on a rockwool substrate by drip irrigation simultaneously with **MYCOSTOP** and **LALSTOP® G46 WG**.

For more information on compatibility, contact the manufacturer and distributors: [www.verdera.fi](http://www.verdera.fi)

Always use plant protection products in a safe manner, in accordance with the label instructions.



LALLEMAND PLANT CARE